arXiv:1601.05180v4 [math.NT] 21 Apr 2016

# On the divisibility of class numbers and discriminants of imaginary quadratic fields

Meng Fai Lim*

**Abstract**

Let $n$ be a squarefree positive odd integer. We will prove that there exist infinitely many imaginary quadratic number fields with discriminant divisible by $n$ and—at the same time—having an element of order $n$ in the class group. We then apply our result to prove that for a given squarefree positive odd integer $n$ there exist infinitely many $N$ such that $n$ divides both $N$ and $r(N)$, where $r(N)$ is the representation number of $N$ as sums of three squares.

Keywords and Phrases: Class number, discriminant, sums of three squares.

Mathematics Subject Classification 2010: 11R11, 11R29, 11R70.

## 1 Introduction

The class group of a number field is certainly one of the most important arithmetic objects in number theory. Even in the case of an imaginary quadratic field there is a great amount of research on the divisibility of the class number. In particular, for a given positive integer $n > 1$, it was shown by many authors that there exist infinitely many imaginary quadratic number fields each with class number divisible by $n$ (see [1, 2, 4, 6, 8, 9, 10, 11, 14, 15, 16, 17, 18, 20, 21, 22, 23, 25, 26]). Recently, there have been studies on the divisibility of class numbers of imaginary quadratic fields under certain constraints on the discriminant. Pertaining to this problem, Byeon and Lee [3], and Pekin [24] have shown that for a given $g \geq 1$, there are infinitely many imaginary quadratic number fields whose discriminant has only two prime factors and whose class number is divisible by $2g$.

In this paper we consider the divisibility of the class numbers under the constraint of the divisibility of the discriminants. More precisely, for a given squarefree odd integer $n > 1$, we will show that there are infinitely many imaginary quadratic number fields $K$ whose class number and discriminant are both divisible by $n$.

**Theorem 1.1.** *Let $n$ be a squarefree odd integer $> 1$. For each congruence class 5 mod 8, 2 mod 4 and 3 mod 4, there exist infinitely many squarefree integers $d < 0$ such that $\gcd(d, h(d))$ is divisible by $n$. Here $h(d)$ is the class number of $\mathbb{Q}(\sqrt{d})$.*

*School of Mathematics and Statistics & Hubei Key Laboratory of Mathematical Sciences, Central China Normal University, Wuhan, 430079, P.R.China. E-mail: limmf@mail.ccnu.edu.cn

**Remark 1.2.** The squarefree condition on $n$ is imposed by our requirement that $n$ divides the discriminant of the imaginary quadratic number field which is in turn squarefree (up to a power of 2). A weaker version of Theorem 1.1 is also proven in [19].

We now give a brief description of the layout of the paper. In Section 2, we recall an algebraic result of Karoubi-Lambre which will be used subsequently in the paper. The proof of Theorem 1.1 will be given in Section 3. In Section 4, we apply Theorem 1.1 to obtain divisibility results on representation number of $N$ as sums of three squares. We also apply Theorem 1.1 to obtain some result on the 3-rank of class groups of real quadratic fields in Section 5. Finally, we give some numerical examples in Section 6 illustrating the results of the paper.

**Acknowledgments.** Some of the material presented in this paper is part of the author's Master thesis [19]. The author would like to thank his supervisor A. Jon Berrick for his advice and encouragement during then. The author would also like to thank Manfred Kolster for his interest and his many valuable comments, which improved the presentation of the paper. The author is also grateful of Peter Cho and Azizul Hoque for their interests and comments on various aspects of related works in the literature. The author is supported by the National Natural Science Foundation of China under the Research Fund for International Young Scientists (Grant No: 11550110172).

## 2 A result of Karoubi and Lambre

We now state the following result of Karoubi and Lambre [12, Proposition 2.8], which our results will depend on. For the proof which relies on an algebraic $K$-theory argument, we refer to the original paper (see also [13] or [19]).

**Theorem 2.1.** *Let $K$ be an imaginary quadratic field with discriminant $D$. Let $n$ be an odd divisor of $D$. Suppose that there exists a triple $(a, b, c) \in \mathbb{Z}^3$ which satisfies all of the following properties*:

(*i*) $(a, b) = (c, n) = 1$.

(*ii*) $b \neq \pm 1$.

(*iii*) $a^2 - 4b^n = c^2 D$.

*Then the class group of $K$ contains an element of order $n$.*

## 3 Proof of Theorem 1.1

In this section, we will prove our main result. As a start, we record the following lemma.

**Lemma 3.1.** *Let $f(X) \in \mathbb{Z}[X]$ be a nonconstant polynomial. Then there exist infinitely many primes $l$ such that the congruence $f(X) \equiv 0$ mod $l$ is solvable.*

*Proof.* If $f(0) = 0$, then this is clearly solvable for all primes. Therefore, we may assume that $f(0) = m \neq 0$. Since $f$ is nonconstant, we can find some $t \in \mathbb{Z}$ such that $|f(t)| > 2$. In particular, there exists at least one prime such that the congruence is solvable. Now suppose that $l_1, ..., l_r$ are primes such that the congruence $f(X) \equiv 0 \bmod l_i$ is solvable. Choose $k$ sufficiently large so that

$$|f((l_1 \cdots l_r)^k m)| > |m|.$$

It then follows that $f((l_1 \cdots l_r)^k m) = c(Nl_1 \cdots l_r + 1)$ for some nonzero integer $N$. Since $N$ is nonzero, there exists a prime divisor $q$ of $Nl_1 \cdots l_r + 1$, which is clearly not equal to any of the $l_i$'s. Hence the lemma is proven. □

We are now in the position to prove Theorem 1.1. For notational convenience, we will write $(s, t)$ for the greatest common divisor of $s$ and $t$.

*Proof of Theorem 1.1.* Since $n$ is assumed to be squarefree and odd, we may write $n = p_1 p_2 \cdots p_r$, where each $p_i$ is an odd prime. Let $a$ be a fixed integer which satisfies all the following properties.

(1) $a \equiv 2 \bmod n$;

(2) $a \not\equiv 2 \bmod p_i^2$ for every $i$;

(3) $(a, 2n) = 1$.

It follows from this choice of $a$ that for each $i$, $X = 1$ is a solution to the congruence $4X^n - a^2 \equiv 0 \bmod p_i$, and that $4 - a^2 = (2 + a)(2 - a) \not\equiv 0 \bmod p_i^2$. Let $T$ be a finite set of odd primes which contains every $p_i$, and has the property that for every other $l$ in $T$, the congruence $4X^n - a^2 \equiv 0 \bmod l$ is solvable with $(l, a) = 1$. Note that the set $T$ can be chosen to be arbitrarily large by Lemma 3.1. We then choose an integer $b > 1$ which satisfies all the following properties.

(1) $4b^n \equiv a^2 \bmod l$ and $4b^n \not\equiv a^2 \bmod l^2$ for every $l \in T$;

(2) $(a, b) = 1$;

(3) $b \equiv 1 \bmod 2$;

(4) $4b^n - a^2 > 0$.

The existence of $b$ follows by an application of the Chinese Remainder Theorem. Since $4b^n - a^2 > 0$ by our choice of $b$, we may write $a^2 - 4b^n = c^2 d$ where $d$ is squarefree and $< 0$. For each $l \in T$, $l$ does not divide $c$, else we have $l^2$ dividing $c^2$ and hence $4b^n - a^2$ contradicting our construction. In particular, we have that $n$ divides $d$ and $(n, c) = 1$. Also, since $a$ is odd (by our initial construction), so is $c$. We now verify that $d \equiv 5 \bmod 8$. To see this, we note that

$$d \equiv c^2 d = a^2 - 4b^n \equiv 1 - 4 \equiv 5 \bmod 8,$$

where the first and third congruences follow from the fact that $a$, $b$ and $c$ are odd. Therefore, the discriminant of the field $K = \mathbb{Q}(\sqrt{d})$ is $d \equiv 5 \bmod 8$. One can check easily that it follows from our construction that the triple $(a, b, c)$ satisfies all the hypotheses in Theorem 2.1, and hence $K = \mathbb{Q}(\sqrt{d})$ contains an element of order $n$. Since the set $T$ can be chosen to be arbitrarily large by Lemma 3.1, it

follows that there exist infinitely many squarefree integers $d < 0$ such that $d \equiv 5 \bmod 8$ and $\gcd(d, h(d))$ is divisible by $n$.

We now show that there exist infinitely many squarefree integers $d < 0$ such that $d \equiv 2 \bmod 4$ and $\gcd(d, h(d))$ is divisible by $n$. Choose an integer $a$ which satisfies all the following properties.

(1) $a \equiv 2 \bmod n$;

(2) $a \not\equiv 2 \bmod p_i^2$ for every $i$;

(3) $a \equiv 2 \bmod 4$.

Let $T$ be a finite set of odd primes which contains every $p_i$, and has the property that for every other $l$ in $T$, the congruence $4X^n - a^2 \equiv 0 \bmod l$ is solvable with $(l, a) = 1$. We then choose an integer $b > 1$ which satisfies all the following properties.

(1) $4b^n \equiv a^2 \bmod l$ and $4b^n \not\equiv a^2 \bmod l^2$ for every $l \in T$;

(2) $(a, b) = 1$;

(3) $b \equiv 3 \bmod 4$;

(4) $4b^n - a^2 > 0$.

We claim that $8 \mid 4b^n - a^2$ and $16 \nmid 4b^n - a^2$. To see this, note that since $a \equiv 2 \bmod 4$, we have $a^2 \equiv 4 \bmod 16$. Since $b = 3 \bmod 4$, we also have $4b^2 \equiv 4 \bmod 16$. Combining these observations with the fact that $n$ is odd, we obtain $4b^n \equiv 4b \equiv 12 \bmod 16$. It then follows that

$$a^2 - 4b^n \equiv 4 - 12 \equiv 8 \bmod 16.$$

Hence we may write $a^2 - 4b^n = 4c^2 d$, where $d$ is squarefree $< 0$, $d \equiv 2 \bmod 4$ and $(2, c) = 1$. By construction, it follows that $n \mid d$ and that the triple $(a, b, c)$ satisfies all the hypotheses in Theorem 2.1 for the field $K = \mathbb{Q}(\sqrt{d})$ (noting that the disciminant of $K$ is $4d$). Hence the class group of $K = \mathbb{Q}(\sqrt{d})$ contains an element of order $n$. Since the set $T$ can be chosen arbitrarily large by Lemma 3.1, it follows that there exist infinitely many squarefree integers $d < 0$ such that $d \equiv 2 \bmod 4$ and $\gcd(d, h(d))$ is divisible by $n$.

Finally, we will prove that there exist infinitely many squarefree integers $d < 0$ such that $d \equiv 3 \bmod 4$ and $\gcd(d, h(d))$ is divisible by $n$. Here the integer $a$ is chosen so that it satisfies all the following properties.

(1) $a \equiv 2 \bmod n$;

(2) $a \not\equiv 2 \bmod p_i^2$ for every $i$;

(3) $a \equiv 0 \bmod 4$.

As before, denote by $T$ a finite set of odd primes which contains every $p_i$, and that for every other $l$ in $T$, the congruence $4X^n - a^2 \equiv 0 \bmod l$ is solvable with $(l, a) = 1$. The integer $b > 1$ is then chosen so that it satisfies all the following properties.

(1) $4b^n \equiv a^2 \bmod l$ and $4b^n \neq a^2 \bmod l^2$ for every $l \in T$;

(2) $(a, b) = 1$;

(3) $b \equiv 1 \bmod 4$;

(4) $4b^n - a^2 > 0$.

We claim that $4 \mid 4b^n - a^2$ and $8 \nmid 4b^n - a^2$. To see this, we first note that since $a \equiv 0 \bmod 4$, we have $a^2 \equiv 0 \bmod 8$. Also, since $b \equiv 1 \bmod 4$, we have $4b^2 \equiv 4 \bmod 8$ which in turn implies that $4b^n \equiv 4b \equiv 4 \bmod 8$ (the first congruence follows from the fact that $n$ is odd). Hence we have that

$$a^2 - 4b^n \equiv 0 - 4 \equiv 4 \bmod 8.$$

Therefore, we may write $a^2 - 4b^n = 4c^2 d$, where $d$ is squarefree $< 0$ and $(2, cd) = 1$. We shall show that $d \equiv 3 \bmod 4$. Suppose not, then we have $d \equiv 1 \bmod 4$ (since $d$ is odd). Now since $c$ is odd, we therefore have $4c^2 \equiv 4 \bmod 16$ which in turn implies that $4c^2 d \equiv 4d \equiv 4 \bmod 16$. Here the last equality follows from our supposition that $d \equiv 1 \bmod 4$. On the other hand, since $a \equiv 0 \bmod 4$ and $b \equiv 1 \bmod 4$, we have $a^2 - 4b^n \equiv -4 \bmod 16$. Combining the two equations, we have $-4 \equiv 4 \bmod 16$, which is a contradiction. Hence we must have $d \equiv 3 \bmod 4$.

By construction, it follows that $n \mid d$ and the triple $(a, b, c)$ satisfies all the hypotheses in Theorem 2.1 for the field $K = \mathbb{Q}(\sqrt{d})$ (noting that the disciminant of $K$ is $D = 4d$). Hence the class group of $K = \mathbb{Q}(\sqrt{d})$ contains an element of order $n$. The proof of the theorem is now completed. □

# 4 Sum of three squares

The class numbers of imaginary quadratic fields are intimately related to the representation numbers of integers as sum of three squares. For a positive integer $N$, we denote by $r(N)$ the number of representations of $N$ as a sum of 3 squares. A classical theorem of Gauss (for instance, see [7, Chap. 4 §8]) asserts that

$$r(N) = \begin{cases} 12H(4N) & \text{if } N \equiv 1, 2 \bmod 4, \\ 24H(N) & \text{if } N \equiv 3 \bmod 8, \\ r(N/4) & \text{if } N \equiv 0 \bmod 4, \\ 0 & \text{if } N \equiv 7 \bmod 8. \end{cases}$$

Here $H(N)$ is the Hurwitz-Kronecker class number (see [5, Definition 5.3.6]). In fact, if $-N = c^2 D$, where $D$ is a negative discriminant $< 4$, we have the following expression (cf. [5, Lemma 5.3.7])

$$H(N) = h(D) \sum_{i \mid c} \mu(i) \Big(\frac{D}{i}\Big) \sigma_1\Big(\frac{c}{i}\Big),$$

where $\mu(i)$ is the Möbius function and $\sigma_1(i)$ is the sum of all positive divisors of $i$.

We can now prove the following which is concerned with the divisibility of $\gcd(n, r(N))$.

**Theorem 4.1.** *Let $n$ be a squarefree odd integer $> 1$. Then for each congruence class 3 mod 8, 2 mod 4 and 1 mod 4, there exist infinitely many squarefree integers $N$ such that $\gcd(N, r(N))$ is divisible by $n$.*

**Remark 4.2.** In [4, Theorem 2], Cho has also established the infinitude of $N$ in certain arithmetic progressions whose $r(N)$ is divisible by $n$. However, his results do not have the constraint on the divisibility of $N$.

We now prove Theorem 4.1.

*Proof of Theorem 4.1.* We first consider the situation when $N \equiv 3 \bmod 8$. Let $K = \mathbb{Q}(\sqrt{d})$ be an imaginary quadratic field, where $d \equiv 5 \bmod 8$ and $\gcd(d, h(d))$ is divisible by $n$. Set $N = -d$. Then $n$ divides $N$ and $N \equiv 3 \bmod 8$. It then follows that

$$r(N) = 24H(N) = 24h(d).$$

This in turn implies that $n$ divides $r(N)$. Since Theorem 1.1 guarantees us that there is an infinite family of quadratic fields to choose, we have our result. The proofs of the remaining cases can be dealt similarly. □

**Remark 4.3.** By a careful examination of our proof, we see that we also prove that for each congruence class 2 mod 4 and 1 mod 4, there exist infinitely many squarefree integers $N$ such that $r(N)$ is divisible by $12n$, and for the congruence class 3 mod 8, there exist infinitely squarefree integers $N$ such that $r(N)$ is divisible by $24n$.

# 5 Real quadratic fields

It would be of interest to establish results analogous to that in Theorem 1.1 for the real quadratic fields. At present, we can only prove results in this direction for the 3-rank of the ideal class groups. Notice that unlike the imaginary situation, our results in the real situation obtain a family of real quadratic fields whose discriminant is *not* divisible by 3.

**Theorem 5.1.** *Let $n$ be a squarefree odd integer $> 1$ which is coprime to 3. For each congruence class 1 mod 8, 2 mod 4 and 3 mod 4, there exist infinitely many squarefree $d > 0$ such that $d$ is divisible by $n$ but not divisible by 3, and such that the 3-rank of the class group of the real number field $\mathbb{Q}(\sqrt{d})$ is at least 2.*

*Proof.* Let $n = 3$. By Theorem 1.1, there exists infinitely many imaginary quadratic fields $K = \mathbb{Q}(\sqrt{d'})$ with $d' \equiv 5 \bmod 8$ and such that $3 \mid d'$ and $3 \mid h(d')$. By [27, Theorem 10.10], the latter implies the 3-rank of the ideal class group of the real quadratic field $\mathbb{Q}(\sqrt{d})$ is at least 2, where $d := -d'/3$. It remains to show that $d \equiv 1 \bmod 8$. Note that since $d' = 5 \bmod 8$ and $3 \mid d'$, we have $d' \equiv 21 \bmod 24$. The required congruence is now immediate from this. The discriminant of $\mathbb{Q}(\sqrt{d})$ is precisely $d = -d'/3$ which is not divisible by 3. The proofs for the other congruence classes are similar. □

# 6 Numerical examples

In this section, we present some numerical examples to illustrate our results.

$\underline{n = 3}$

$5^2 - 4(7)^3 = -1347 = -3 \cdot 449$. Note that $-1347 \equiv 5 \bmod 8$. Therefore, the class number and the discriminant of $\mathbb{Q}(\sqrt{-1347})$ are divisible by 3. It then follows that $r(1347)$ is divisible by 72 and the 3-rank of the class group of $\mathbb{Q}(\sqrt{449})$ is at least 2.

$5^2 - 4(43)^3 = -318003 = -3 \cdot 7 \cdot 19 \cdot 797$. Note that $-318003 \equiv 5 \bmod 8$. Therefore, the class number and the discriminant of $\mathbb{Q}(\sqrt{-318003})$ are divisible by 3. It then follows that $r(318003)$ is divisible by 72 and the 3-rank of the class group of $\mathbb{Q}(\sqrt{106001})$ is at least 2.

$14^2 - 4(55)^3 = -665304 = -8 \cdot 83163 = -2^3 \cdot 3 \cdot 19 \cdot 1459$. Note that $-166326 = -2 \cdot 83163 \equiv 2 \bmod 4$. Therefore, the class number and the discriminant of $\mathbb{Q}(\sqrt{-166326})$ are divisible by 3, $r(166326)$ is divisible by 36 and the 3-rank of the class group of $\mathbb{Q}(\sqrt{55442})$ is at least 2.

$\underline{n = 5}$

$16^2 - 4(29)^5 = -2^2 \cdot 20511085 = -2^2 \cdot 5 \cdot 7 \cdot 151 \cdot 3881$. Note that $-20511085 \equiv 3 \bmod 4$. Therefore, the class number and the discriminant of $\mathbb{Q}(\sqrt{-20511085})$ are divisible by 5, and $r(20511085)$ is divisible by 60.

$\underline{n = 7}$

$9^2 - 4(8)^7 = -8388527 = -7 \cdot 1198361$. Note that $-8388527 \equiv 5 \bmod 8$. Therefore, the class number and the discriminant of $\mathbb{Q}(\sqrt{-8388527})$ are divisible by 5, and $r(8388527)$ is divisible by 168.

$\underline{n = 15 = 3 \cdot 5}$

$$490912124320572 - 4(61)^{15} = -90813862366184355 = -3 \cdot 5 \cdot 73093973 \cdot 82828409.$$

Note that 73093973 and 82828409 are primes, and $-90813862366184355 \equiv 5 \bmod 8$. Therefore, the class number and the discriminant of $\mathbb{Q}(\sqrt{-90813862366184355})$ are divisible by 15, and $r(90813862366184355)$ is divisible by 360. Furthermore, the 3-rank of the class group of $\mathbb{Q}(\sqrt{30271287455394785})$ is at least 2.

$$490912123905322 - 4(61)^{15} = -41678390531241925 80 = -2^2 \cdot 3 \cdot 5 \cdot 69463984218736543.$$

Note that 69463984218736543 is a prime and $-1041959763281048145 \equiv 3 \bmod 4$. Therefore, the class number and the discriminant of $\mathbb{Q}(\sqrt{-1041959763281048145})$ are divisible by 15, and $r(1041959763281048145)$ is divisible by 180. Furthermore, the 3-rank of the class group of $\mathbb{Q}(\sqrt{347319921093682715})$ is at least 2.

# References


[1] N. C. Ankeny and S. Chowla, On the divisibility of the class number of quadratic fields, *Pacific J. Math.* 5 (1955) 321-324.

[2] D. Byeon, Divisibility properties of class numbers, *Trends in Mathematics*, Information Center for Mathematical Sciences, Vol 4 No. 1, June 2001, pp 26-30.

[3] D. Byeon and S. Lee, Divisibility of class numbers of imaginary quadratic fields whose discriminant has only two prime factors, *Proc. Japan Acad.* 84 (2008) 8-10.

[4] P. J. Cho, Sum of three squares and class numbers of imaginary quadratic fields, *Proc. Japan Acad. Ser. A Math. Sci.* 87(6) (2011) 91-94.

[5] H. Cohen, *A Course in Computational Algebraic Number Theory*, Grad. Texts in Math. **138**, Springer-Verlag, New York, 1993.

[6] B. Gross and D. Rohrlich, Some results on the Mordell-Weil group of the Jacobian of the Fermat curve, *Invent. Math* 44 (1978) 201-224.

[7] E. Grosswald, *Representations of Integers as Sums of Squares*, Springer-Verlag, New York, 1985.

[8] A. Hoque and H. K. Saikia, A family of imaginary quadratic fields whose class numbers are multiples of three, *J. Taibah. Univ. Sci.* 9 (2015) 399-402.

[9] A. Hoque and H. K. Saikia, A note on quadratic fields whose class numbers are divisible by 3, *SeMA. J.* 73 (2016) 1-5.

[10] P. Humbert, Sur les nombres de classes de certains corps quadratiques, *Comment. Math. Helv.* 12 (1940) 233-245.

[11] P. Humbert, Note relative á l'árticle: Sur les nombres de classes de certains corps quadratiques, *Comment. Math. Helv.* 13 (1940) 67.

[12] M. Karoubi and T. Lambre, Quelques classes caractéristiques en théorie des nombres, *J. reine angew. Math.* 543 (2002) 169-186.

[13] M. Karoubi and T. Lambre, *Quelques classes caractéristiques en théorie des nombres*, School and Conference on Algebraic K-Theory and its Applications, 8-26 July 2002, The Abdus Salam International Centre for Theoretical Physics, SMR 1418/16.

[14] Y. Kishi, A new family of imaginary quadratic fields whose class number is divisible by five, *J. Number Theory* 128 (2008) 2450-2458.

[15] Y. Kishi, Note on the divisibility of the class number of certain imaginary quadratic fields, *Glasg. Math. J.* 51(1) (2009) 187-191.

[16] Y. Kishi, Note on the divisibility of the class number of certain imaginary quadratic fields-corrgendum, *Glasg. Math. J.* 52(1) (2010) 207-208.

[17] Y. Kishi, On the ideal class group of certain quadratic fields, *Glasg. Math. J.* 52(3) (2010) 575-581.

[18] S. -N. Kuroda, On the class number of imaginary quadratic number fields, *Proc. Japan Acad.* 40 (1964) 365-367.

[19] M. F. Lim, Some characteristic classes in number theory, Master thesis, National University of Singapore, 2005.

[20] S. Louboutin, On the divisibility of the class number of imaginary quadratic number fields, *Proc. Amer. Math. Soc.* 137 (2009), 4025-4028.

[21] R. Mollin, On class numbers of quadratic extensions of algebraic number fields, *Proc. Japan Acad.* 62 (1986) 33-36.

[22] M. R. Murty, Exponents of class groups of quadratic number fields, Topics in Number Theory (University Park, PA, 1997) Kluwer Acad. Publ., Dordrecht (1999), 229-239.

[23] T. Nagell, Über die Klassenzahl imaginär-quadratischer Zahlköper, *Abh. Math. Sem. Univ. Hamburg* 1 (1922) 140-150.

[24] A. Pekin, Divisibility criteria for class numbers of imaginary quadratic fields whose discriminant has only two prime factors, *Abstr. Appl. Anal.* 2012, Art. ID 570154, 4 pp.

[25] K. Soundararajan, Divisibility of class numbers of imaginary quadratic fields, *J. London Math. Soc. (2)* 61(3) (2003) 681-690.

[26] T. Uehara, On class numbers of imaginary quadratic and quartic fields, *Archiv. der Math.* 41(3) (1983) 256-260.

[27] L. C. Washington, *Introduction to Cyclotomic Fields*, 2nd edn., Grad. Texts in Math. **83**, Springer-Verlag, New York, 1997.